

TORAKS 2021

11. kongres Hrvatskog torakalnog društva 11th Congress of the Croatian Thoracic Society 19.-22. svibnja | May Virtualni kongres | Virtual congress

NT-PROBNP AS A TOOL TO STRATIFY SEVERITY OF PULMONARY HYPERTENSION IN CHRONIC OBSTRUCTIVE PULMONARY DISEASE

PAVLIŠA G.^{1, 2}, Ljubičić L.¹, Hećimović A.¹, Basara L.¹, Janković Makek M.^{1, 2}, Džubur F.^{1, 2}, Glodić G.¹, Puretić H.¹, Tolić E.², Samaržija M.^{1, 2}

¹ University Hospital Center Zagreb, Zagreb, Croatia Clinic for Respiratory Diseases Jordanovac

² University of Zagreb , Zagreb, Croatia School of Medicine

Objective:

Pulmonary hypertension (PH) is a common complication of chronic obstructive pulmonary disease (COPD) that has an important implication on its prognosis. PH carries the risk of more frequent hospitalizations and shorter survival. However, it is often overlooked due to the difficulty in diagnosing. Echocardiographic assessment of PH may be subject to errors because of poor visualization of the heart due to emphysema, while right heart catheterization (RHC) is an invasive method. The aim of our study was to examine whether some routinely evaluated noninvasive parameter can predict which patient has pulmonary hypertension and requires invasive diagnostic procedure.



Methods: The study included 59 COPD patients (38 men, 21 women, age 60.9 ± 11.1), with RHC confirmed PH. The following tests were performed: complete blood count (CBC), NT- pro-BNP, troponin T, arterial blood tests, pulmonary functional test.

Results: The results of descriptive statistics are presented as mean +/- SD. The parameters obtained by the right heartcatheterization were as follows: mean right atrium *pressure* (mRA) 10,7 +/- 5,33 mmHg, mean pulmonary artery pressure (mPAP) 46,68 +/- 13,66 mmHg, wedge 12,87 +/-13,66 mmHg, cardiac output (CO) 5,34 +/- 1,21 L/min, cardac index (CI) 2,8 +/- 0,6 L/min/m². The prevalence of mild pulmonary hypertension was 17% (10/59), moderate 36% (21/59) and severe 47% (28/59). Pulmonary function parameters were: FVC 64,40 +/- 21,29 % predicted, FEV1 46,08 +/- 20,77 % predicted, FEV1 / FVC 0,54 +/- 0,12, DICO 37,97 +/- 21,26 % predicted. Laboratory parameters were: erythrocytes 5,13 +/- 0.64 x 1012/L, hemoglobin 150,6 +/- 20,0 g/L, hematocrit 0,46 +/- 0.6, troponin T 14,05 +/- 20,04 ng/ml, NT-proBNP 1806 +/- 2191,69 ng/L, pH 7,43 +/-0,38, PaO2 65,78 +/- 9,3 mmHg, PaCO2 44,16 +/- 10,58 mmHg, PaO2/FiO2 289,5 +/-55,84. A significant correlation was found between mPAP and NT-proBNP (R = 0.311, p = .037).

Conclusion: NT-proBNP is released as a result of atrial and ventricular wall stretching. In addition being a marker of left heart failure, its elevated value may also indicate a load on the right heart. There are few studies on its value in identification of pulmonary hypertension in COPD patients, and their results indicate that NT-proBNP has relatively good sensitivity and specificity. The results of our study also suggest that NT-proBNP can predict the severity of pulmonary hypertension in COPD. Further research is needed to assess its value as a screening tool for noninvasive detection of PH in COPD patients.